

Quasi-localized states in disordered metals and non-analyticity of the level curvature distribution function.

V. E. Kravtsov^{1,2} and I. V. Yurkevich¹.

¹ *International Center for Theoretical Physics, P.O. Box 586, 34100 Trieste, Italy*

² *Landau Institute for Theoretical Physics, Kosygina str. 2, 117940 Moscow, Russia*
(February 1, 2008)

It is shown that the quasi-localized states in weakly disordered systems can lead to the non-analytical distribution of level curvatures. In 2D systems the distribution function $P(K)$ has a branching point at $K = 0$. In quasi-1D systems the non-analyticity at $K = 0$ is very weak, and in 3D metals it is absent at all. Such a behavior confirms the conjecture that the branching at $K = 0$ is due to the multi-fractality of wave functions and thus is a generic feature of all critical eigenstates. The relationship between the branching power and the multi-fractality exponent $\eta(2)$ is derived.

PACS numbers: 71.25.-s, 72.15.Rn, 05.45.+b

The problem of energy level statistics in complex quantum systems has recently attracted much attention. It is a key problem for the description of mesoscopic fluctuations in small but macroscopic disordered electronic systems at low temperatures¹. It is also of great fundamental importance, since level statistics is a proper language to describe the signature of chaos in a quantum system whose classical counterpart is chaotic.

The most of investigations in the field has been aimed to describe the spectral statistics themselves independently of the nature of wave functions. Nonetheless, it is the space structure of wave functions that determines the properties of spectral statistics. In systems where the wave functions are extended and essentially structureless (e.g. in good metals) the predictions of the random matrix theory (RMT)² is proved to work well³. The corresponding Wigner-Dyson (WD) spectral statistics is characterized by the level repulsion. On the other hand, in the region of localized states, energy levels are almost uncorrelated obeying the Poisson statistics. The spectral statistics at the mobility edge^{4,5} proves to be universal but different from both the Poisson and the WD statistics. Since the wave functions at the mobility edge are still extended, the energy levels repel each other at small energy separations ω of the order of the mean level spacing Δ , the correlations decaying as a power law⁵ for $\omega \gg \Delta$. However, the typical fluctuation of the number of levels δN in an energy window of the width $E = \bar{N}\Delta \gg \Delta$ is much bigger than the RMT prediction and behaves in a quasi-Poisson way^{6,7}:

$$\langle (\delta N)^2 \rangle = \chi \bar{N} \gg \ln \bar{N}. \quad (1)$$

Such a complex behavior of the critical spectral statistics is due to the complex statistics of the critical wave func-

tions. A typical wave function at the mobility edge can be considered as a superposition of peaks with the broad distribution of heights and widths. The corresponding quantum-mechanical probability density $|\Psi(\mathbf{r})|^2$ is sparse and is not negligible only in a small fraction of volume that scales with the system size L as L^D , $D = d - \eta$ being the fractal dimensionality in the d -dimensional space. However, two wave functions of different energies $|E - E'| \gg \Delta$ overlap well⁸ with each other since they are strongly correlated in a given point:

$$\langle |\Psi_E(\mathbf{r})|^p |\Psi_{E'}(\mathbf{r})|^p \rangle \propto |E - E'|^{-\eta(p)/d}. \quad (2)$$

Such slowly decaying correlations are typical of the critical wave functions. Both in good metal and in the insulator wave functions of different energies are locally uncorrelated. The statistics of wave functions expressed by Eq.(2) exhibits *multi-fractality*⁹, since the fractal dimensionality $D(p) = d - \eta(p)$ depends on the power p . The main effect of multi-fractality is described by the exponent $\eta(2) = \eta$. In particular the level compressibility χ in Eq.(1) has been found⁷ to depend on η as follows:

$$\chi = \frac{\eta}{2d}. \quad (3)$$

Multi-fractality of wave functions is typical not only for critical states at the mobility edge. Weak multi-fractality holds¹⁰⁻¹² for two-dimensional disordered systems in the regime of weak localization where the system size L is much less than the localization radius R_0 .

In general, multi-fractality can be considered as a result of proliferation of the *quasi-localized* states. The latter notion^{13,14} means rare states in disordered systems where $|\Psi(\mathbf{r})|^2$ has a sharp peak on top of the structureless extended background (in 3D metals) or localized states with anomalously small localization radius (for $d \leq 2$). The existence of such states exhibits itself in the slow relaxation of current first discovered in Ref. [11] using the complicated renormalization-group analysis of an extended nonlinear sigma-model in $2 + \epsilon$ dimensions.

Very recently the problem of current relaxation in disordered conductors has been reconsidered^{14,15} by an elegant instanton approximation¹⁴ applied to supersymmetric version of the nonlinear sigma-model³. In these works the main result of the previous consideration¹¹ has been confirmed for 2D systems. However, the new method was able to describe some unknown regimes of current relaxation and to set correct limits of validity for the regimes found earlier. Later the same idea¹⁴

has been applied¹² to find directly the distribution of $|\Psi(\mathbf{r})|^2$ and the distribution of local densities of states¹⁶. Thus the existence of the quasi-localized states has been proved and the corresponding configuration of the random impurity potential has been found^{14,17}.

The question we address in this Letter is whether and how the quasi-localized states and multi-fractality manifest itself in level statistics other than the level number variance. The natural candidate is the statistics of level curvatures K_n first introduced by Thouless¹⁸:

$$K_n = \frac{1}{\Delta} \left. \frac{d^2 E_n(\phi)}{d\phi^2} \right|_{\phi \rightarrow 0}. \quad (4)$$

Here $E_n(\phi)$ is an energy level depending on the twist of phase in the boundary condition $\Psi(\mathbf{r}_\perp, x=0) = \Psi(\mathbf{r}_\perp, x=L)e^{i\phi}$. Such a phase shift arises if an external magnetic flux threads a sample of the toroidal form.

The main idea of Ref. [18] is that it is possible to distinguish between the localized and extended states studying the sensitivity of spectrum to the twist of phase in the boundary conditions. This sensitivity is maximal for extended states and negligible for the localized ones. It is clear that the existence of the quasi-localized states should lead to an enhancement in the probability to find a state with a small value of the random quantity K_n . Thus the quantity of interest is the deviation of the distribution function $P(K)$ from the form^{19,20} that follows from the random matrix theory.

This deviation has been recently studied numerically²¹ and analytically^{20,22} using an expansion in powers of the inverse dimensionless conductance g^{-1} . The main result of the numerical simulations²¹ on the mobility edge of the 3D Anderson model is that the deviation $\delta P(K) = P(K) - P_{WD}(K) \propto |K|^{2-\alpha}$ has a branching point at $|K| = 0$ with $\alpha \approx 0.4$. Although the analytical result²² describes correctly the sign and shape of the deviation $\delta P(K)$, it is unable to explain the branching at $K = 0$ which should be a non-perturbative in $1/g$ effect. It has been also conjectured in²¹ that the branching is due to the multi-fractality of critical wave functions and thus the new critical exponent α should be related with the multi-fractality exponent η .

In this Letter we will show using the instanton approximation similar to that used in¹⁴ that the level curvature distribution $P(K)$ has, indeed, a branching point at $K = 0$ in 2D systems under the weak-localization condition $R_0 \gg L$. In 3D metals and quasi-1D systems with $R_0 \gg L$ the distribution function deviates from the WD form but all its derivatives are finite at $K = 0$. This result favors the statement²¹ that the branching in $P(K)$ at $K = 0$ and multi-fractality are related with each other, since for good conductors the multi-fractality is also present only for $d=2$.

We have found an explicit relationship between α and η that we believe is valid at the mobility edge near the Anderson transition too:

$$\alpha = 3 - 4/\eta, \quad (\eta > 4/3). \quad (5)$$

Another relationship can be found using Eq.(3):

$$\alpha = 3 - \frac{2}{d} \chi^{-1}, \quad (2/d > \chi > 2/3d). \quad (6)$$

The first of these equations relates the parametric level statistics $P(K)$ with the multi-fractal statistics, Eq.(2), of critical wave functions. The second one establishes a connection between the parametric and the conventional level statistics at the mobility edge.

The level curvature distribution function $P(K) = \langle \delta(K - K_n) \delta(E - E_n) \rangle$ can be expressed^{20,22} in terms of the supersymmetric nonlinear sigma-model derived in³ from the model of non-interacting electrons in a random Gaussian potential:

$$P(K) = \lim_{\phi \rightarrow 0} \phi^2 \left[\Re \int \mathcal{D}Q A[Q] e^{-F[Q; K, \phi]} \right], \quad (7)$$

where $A[Q]$ is a certain pre-exponent that is irrelevant for the present consideration and

$$F[Q; K, \phi] = \frac{\pi}{8} g \int Str (\partial Q - i\phi \mathbf{n} [Q, \hat{\tau}])^2 d^d r + i \frac{\pi}{8} K \phi^2 \int Str [\Lambda Q] d^d r. \quad (8)$$

Here $g = \Delta^{-1} D/L^2$ is the dimensionless conductance, D is the diffusion coefficient; the unit vector $\mathbf{n}$ is directed along the x -axis. The field $Q(\mathbf{r})$ is a 8×8 super-matrix of the certain symmetry³ that obeys the condition $Q^2 = 1$. The symbol $Str[.]$ stands for the super-trace. The matrices $\hat{\tau} = \sigma_z(1 + \Lambda)/2$ and $\Lambda = diag(1, -1)$, describe breaking of time-reversal symmetry and the symmetry between retarded and advanced Green's functions, respectively. Here and below we use the dimensionless coordinates-ordinates $\mathbf{r} = \{r_i\}$, where $-1/2 < r_i < 1/2$.

One can see that the level curvature K enters Eq.(8) in the same way as the frequency ω in the problem of long-time current relaxation¹⁴. Therefore from the results of Ref.[14] one would expect the instanton approximation to work well for the Fourier-transform $\tilde{P}(\lambda) = \int dK exp[-iK\lambda]$ for $\lambda \gg 1$ rather than for the distribution function $P(K)$ itself.

It is very important that in order to find the distribution function $P(K)$ or $\tilde{P}(\lambda)$ one should do the limit $\phi \rightarrow 0$. In this limit a nonzero contribution to the functional $F[Q; K, \phi \rightarrow 0]$ can arise only from the components in the non-compact boson-boson sector of the field Q that can take arbitrary large values.

Another simplification comes from the fact that integration over Grassman variables requires an expansion of $exp\{-F[Q; K, \phi]\}$ in powers of the Grassman components of the field Q . Therefore such components can only contribute to the pre-exponent A in the expression $\tilde{P}(\lambda) = A e^{-S(\lambda)}$, where the effective action $S(\lambda) \gg 1$. The main, exponential effect is due to the conventional variables that parametrize the matrix field Q . Thus using

the Efetov's parametrization³ for the orthogonal ensemble we have $Q = V^{-1}HV$, where H contains the "non-compact" angles $\theta_{1,2}(\mathbf{r}) \geq 1$ and V contains "phase" variables $\varphi(\mathbf{r}) \in [0, 2\pi]$ and $\chi(\mathbf{r}) \in [0, 2\pi]$. All functions must obey the periodic boundary conditions.

By varying the functional $F[Q; K, \phi] + iK\lambda$ over these functions we find the saddle-point equations to be satisfied by the choice $\partial\chi = 0$, $\theta_1 = \theta_2 = \theta$ and $\varphi(\mathbf{r}) = \phi v(\mathbf{r}) \neq 0$. The equations for the rest two functions read:

$$\partial^2\theta + \phi^2[\kappa - (\partial v - \mathbf{n})^2] \sinh\theta = 0, \quad (9)$$

$$\partial[(\partial v - \mathbf{n})(\cosh\theta - 1)] = 0. \quad (10)$$

The variation over $K = -2ig\kappa$ leads to the self-consistency equation, Eq.(14).

The limit $\phi \rightarrow 0$ is done simply by absorbing ϕ^2 into θ . We introduce $\tilde{\theta} = \theta + \ln\phi^2$. Then in the limit $\phi \rightarrow 0$ we have $\sinh\theta = \cosh\theta = \frac{1}{2}e^{\tilde{\theta}}\phi^{-2} \gg 1$ so that ϕ drops from Eqs.(9),(10).

One can solve Eq.(10) to obtain:

$$(\partial v - \mathbf{n}) = \text{rot}\mathbf{A} e^{-\tilde{\theta}}, \quad (11)$$

where $\mathbf{A}$ is an arbitrary vector function. In what follows we limit ourselves by considering $\text{rot}\mathbf{A} = -\mathbf{n}/N = \text{const}$. Integrating Eq.(11) over space and using the periodic boundary conditions for $v(\mathbf{r})$ we obtain the second self-consistency equation Eq.(13).

All equations take a more symmetric form if we make a shift $\theta = u - \zeta$, where $\sinh\zeta = N\kappa^{-1/2}(\kappa - N^{-2})/2$:

$$\partial^2 u + \gamma^2 \sinh u = 0, \quad (12)$$

$$\frac{1}{N} = \gamma^2 \int e^{-u} d^d r, \quad (13)$$

$$\lambda = \frac{\pi}{8\gamma^2 N^2} \int e^u d^d r, \quad (14)$$

where $\gamma^2 = \sqrt{\kappa}/N$.

Solving these equations with the periodic boundary conditions one finds $u(\mathbf{r}, \lambda)$, $N(\lambda)$ and $\gamma(\lambda)$ that enter the instanton action $S(\lambda) = F[Q; K, \phi \rightarrow 0] + iK\lambda|_{s.p.}$:

$$-\ln \tilde{P}(\lambda) \sim S(\lambda) = \frac{\pi}{4}g \int (\partial u)^2 d^d \rho + 2g\gamma^4 N^2 \lambda. \quad (15)$$

Eqs.(12),(14) are analogous to those derived in^{14,15} for the problem of current relaxation in the orthogonal ensemble. The second self-consistency equation (13) is typical for our problem which is essentially a crossover problem from the orthogonal to the unitary ensemble. Another important difference is that we are looking for the *periodic* solutions to the above equations with the period $a = 1$ rather than the solutions¹⁴ that vanish at the boundary of the sample.

Since $\gamma^2 \propto K^{1/2}$ we expect the saddle point to correspond to $\gamma^2 \ll 1$ at $\lambda \rightarrow \infty$. In this case the nonlinear term in Eq.(12) can be approximated by $\gamma^2 \sinh u \approx \frac{\gamma^2}{2}e^{|u|} \text{sign}(u)$. Thus we come to the Liouville equation.

Let us first consider the case of a quasi-1D sample. Then the proper periodic solution to the Liouville equation takes the form:

$$e^u = 4k^2 \gamma^{-2} \cosh^{-2}(kx), \quad (|x| < 1/4), \quad (16)$$

where $4k^2 = \gamma^2 \cosh^2(k/4)$. In order to obtain the solution $u(x)$ for $1/4 < |x| < 1/2$ one should simply continue the solution, Eq.(16), anti-symmetrically about the points $x = \pm 1/4$. Because of such anti-symmetry we have

$$\int e^{-u} d^d r = \int e^{+u} d^d r. \quad (17)$$

Finally for the case of quasi-1D sample we obtain $N^{-1} = 8k \approx 8 \ln(1/\gamma^4)$, $\gamma^4 = 64\pi k^3/\lambda$, and

$$\tilde{P}(\lambda) = A \exp \left[-\frac{g_1}{2} \ln^2 \lambda \right], \quad (18)$$

where $g_1 = 2\pi g$. This result is *exactly* of the same form as the long tails of current relaxation¹⁴ (in the orthogonal ensemble) if one replaces $t \rightarrow \lambda$. The region of validity $1 \ll \lambda \ll e^{L/l}$ of Eq.(18) is found from the condition of the breakdown of the diffusive regime^{14,15} $|\partial u(\mathbf{r})|_{\max} = L/l$, where l is the elastic scattering length.

In the 2D case we will look for the continuous periodic solution to the Liouville equation with the symmetry of a square. It is positive inside the square Ω with vertices at $(0; \pm 1/2)$ and $(\pm 1/2; 0)$ and is negative in the rest half of the sample, being anti-symmetric with respect to all sides of the square Ω and thus obeying Eq.(17).

The problem of finding such a solution to the Liouville equation can be solved exactly using the conformal transformations of a unit circle onto the square Ω . However, for our purposes it is sufficient to know the solution for $r \ll 1$ where it depends only on r . The solution that leads to the finite action $S(\lambda)$ is given by:

$$e^{u(r)} = 16b(\gamma^2 + br^2)^{-2}, \quad (19)$$

where $b = 4\pi^4$. Using this solution we find $N^{-1} = 16\pi$, $\gamma^4 = 8^3\pi^4/\lambda$, and

$$\tilde{P}(\lambda) = A \lambda^{-2g_2}. \quad (20)$$

where $g_2 = 2\pi^2 g$. Besides the validity condition $1 \ll \lambda \ll (L/l)^4$, this result has again exactly the same functional form as the long tails of current relaxation¹⁴ in 2D systems.

In the 3D metal phase, the saddle-point solution for $\tilde{P}(\lambda)$ that depends only on one parameter, the dimensionless conductance, is absent. The reason is that the 3D Liouville equation does not have a solution that corresponds to a finite action $S(\lambda)$ in the limit $L/l \rightarrow \infty$.

It is remarkable that only in the 2D case the instanton contribution to the Fourier-transform $\tilde{P}(\lambda)$ of $P(K)$ has a power-law asymptotic behavior at $\lambda \gg 1$. In the limit ²³ $L/l \rightarrow \infty$ this behavior is not restricted from above and the distribution function $P(K)$ must have a branching point at $K = 0$. Namely, there should exist a set of critical values of $g_2^{(n)} = \frac{2n+1}{2} = \frac{1}{2}, \frac{3}{2}, \dots$ such as for $\frac{2n-1}{2} < g_2 < \frac{2n+1}{2}$ the derivative $d^{2n}P(K)/dK^{2n} \propto |K|^{-\alpha_n}$ at $|K| \rightarrow 0$. The value of α_n is given by:

$$\alpha_n = (2n + 1) - 2g_2. \quad (21)$$

In particular, for $g_2 < \frac{1}{2}$, the distribution function $P(K)$ should be divergent at $K \rightarrow 0$. For $\frac{1}{2} < g_2 < \frac{3}{2}$ it should behave like $P(K) - P(0) \propto |K|^{2-\alpha}$ where $\alpha = \alpha_1 = 3 - 2g_2$. One can eliminate the dependence on g_2 using the expression ¹² for the multi-fractality exponent $\eta = \frac{2}{g_2}$. Then we arrive at the result, Eq.(5).

In the quasi-1D case the function $\tilde{P}(\lambda)$ falls off fast enough, and all derivatives of $P(K)$ are finite at $K = 0$:

$$P_{2n} \equiv d^{2n}P/dK^{2n}|_{K=0} \propto \exp[(2n + 1)^2/2g_1]. \quad (22)$$

In this case we have much softer non-analyticity at $K = 0$ than that in 2D case. However, the function $P(K)$ is still not regular at $K = 0$, since the Taylor series $\sum_n P_{2n}K^{2n}/(2n)!$ has zero radius of convergence.

Only in the 3D case the quasi-localized states are not that important in the metal phase and the correction $P(K) - P_{WD}(K)$ can be found perturbatively in $1/g$ as it has been done in Ref.[22].

The above treatment does not allow to take into account localization effects explicitly. Thus we are not able to *derive* the level curvature distribution at the mobility edge of 3D disordered systems. However, we believe that the distribution function at the mobility edge is similar to the one in 2D systems, since both cases are characterized by the multi-fractal statistics of wave functions. That is why we believe the relationship, Eq.(5), that we derived for 2D systems in the weak localization regime, to hold also in the critical region near the Anderson transition.

In summary, we have shown that the quasi-localized states and multi-fractality of wave functions can lead to a non-analytical distribution of level curvatures K . The distribution function $P(K)$ is shown to have a branching point at $K = 0$ for 2D systems and a softer non-analyticity in quasi-1D systems in the weak-localization regime. In the 3D metal the distribution function is analytical at $K = 0$. We argue that the branching in $P(K)$ is typical for all critical states where the statistics of eigenfunctions exhibit multi-fractality. This allows to conjecture a general relationship between the branching power and the multi-fractality exponent η .
Acknowledgments We thank C.M.Canali, V.I.Falko, Y.V.Fyodorov, I.V.Lerner, Yu Lu and A.D.Mirlin for stimulating discussions. Support from the RFBR-INTAS grant no. 95-0675 (V.E.K.) is gratefully acknowledged.

-
- [1] *Mesoscopic Phenomena in Solids*, ed. B. L. Altshuler et al, Elsevier, Amsterdam 1991.
 - [2] M. L. Mehta, *Random matrices* (Academic Press, Boston, 1991).
 - [3] K. B. Efetov, *Adv. Phys.* **32**, 53 (1983).
 - [4] B. I. Shklovskii, B. Shapiro, B. R. Sears, P. Lambrianides, and H. B.Shore, *Phys. Rev. B.* **47**, 11487 (1993).
 - [5] V. E. Kravtsov, I. V. Lerner, B. L. Altshuler and A. G. Aronov, *Phys. Rev. Lett.* **72**, 888 (1994); A. G. Aronov, V. E. Kravtsov, and I. V. Lerner, *Phys. Rev. Lett.* **74**, 1174 (1995).
 - [6] B. L. Altshuler, I. Kh. Zharekeshev, S. A. Kotochigova and B. I. Shklovskii, *Zh. Exp. Tior. Fiz.* **94**, 343 (1988) [*Sov. Phys. JETP* **67**, 625 (1988)].
 - [7] J. T. Chalker, V. E. Kravtsov and I. V. Lerner, *Pis'ma Zh. Exp. Teor. Fiz.* **64**, 355 (1996)
 - [8] J. T. Chalker, *Physica A* **167**, 253 (1990); B. Huckenstein and L. Schweitzer, *Phys. Rev. Lett.* **72**, 713 (1994).
 - [9] C. Castellani and L. Peliti, *J. Phys. A* **19**, L429 (1986); W. Pook and M. Janssen, *Z. Phys. B.* **82**, 295 (1991).
 - [10] F. Wegner, *Z. Phys. B* **36**, 209 (1980).
 - [11] B. L. Altshuler, V. E. Kravtsov and I. V. Lerner in Ref.[1], p.449 and references therein.
 - [12] V. I. Falko, K. B. Efetov, *Phys. Rev. B.* **52**, 17413 (1995).
 - [13] V. E. Kravtsov, *Habilitationsschrift*, Heidelberg University 1992 (unpublished).
 - [14] B. A. Muzykantskii and D. E. Khmelnitskii, *Phys. Rev. B.* **51**, 5480 (1995); *JETP Lett.* **62**, 76 (1995).
 - [15] A. D. Mirlin, *Pis'ma Zh. Exp. Teor. Fiz.* **62**, 583 (1995) [*JETP-Lett.* **62**, 603 (1995)].
 - [16] A. D. Mirlin, *Phys. Rev. B.* **53**, 1186 (1996).
 - [17] I. E. Smolyarenko and B. L. Altshuler, *cond.-math/9606181* (unpublished).
 - [18] D. J. Thouless, *Phys. Rep.* **13**, 93 (1974).
 - [19] F. von Oppen, *Phys. Rev. E.* **51**, 2647 (1995).
 - [20] Y. V. Fyodorov and H.- J. Sommers, *Phys. Rev. E.* **51**, R2719 (1995).
 - [21] C. M. Canali, C. Basu, W. Stephan and V. E. Kravtsov, *Phys. Rev. B.* **54**, 1431 (1996).
 - [22] I. V. Yurkevich and V. E. Kravtsov, *cond-math/9609106* (unpublished).
 - [23] The limit $L/l \rightarrow \infty$ should be taken at a constant g by a simultaneous increase in the number of transverse channels.